

4-(1,2,4-Triazol-1-yl)aniline

Hoong-Kun Fun,^{a,†} Ching Kheng Quah,^{a,§}
B. Chandrakantha,^b Arun M. Isloor^c and Prakash Shetty^d

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Chemistry, Manipal Institute of Technology, Manipal 576 104, India, ^cOrganic Chemistry Division, Department of Chemistry, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India, and ^dDepartment of Printing, Manipal Institute of Technology, Manipal 576 104, India

Correspondence e-mail: hkfun@usm.my


Received 10 December 2010; accepted 11 December 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.118; data-to-parameter ratio = 19.6.

In the title compound, $\text{C}_8\text{H}_8\text{N}_4$, the dihedral angle between the triazole ring [maximum deviation = 0.003 (1) Å] and the benzene ring is 34.57 (7)°. In the crystal, molecules are linked into sheets lying parallel to the *ac* plane via intermolecular $\text{N}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds. Aromatic $\pi-\pi$ [centroid-centroid distance = 3.6750 (8) Å] stacking and $\text{N}-\text{H}\cdots\pi$ interactions are also observed.

Related literature

For general background to and the biological activity of triazole derivatives, see: Isloor *et al.* (2000, 2009); Soliman *et al.* (2001); Holla *et al.* (2000); Sunil *et al.* (2009). For bond-length data, see: Allen *et al.* (1987). For a related structure, see: Fun *et al.* (2010).



Experimental

Crystal data

$\text{C}_8\text{H}_8\text{N}_4$ $c = 19.5477$ (5) Å
 $M_r = 160.18$ $\beta = 99.416$ (2)°
 Monoclinic, $P2_1/c$ $V = 788.15$ (3) Å³
 $a = 5.5488$ (1) Å $Z = 4$
 $b = 7.3656$ (2) Å Mo $K\alpha$ radiation

† Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5525-2009.

$\mu = 0.09$ mm⁻¹
 $T = 296$ K

0.50 × 0.42 × 0.14 mm

Data collection

Bruker SMART APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2009)
 $T_{\min} = 0.957$, $T_{\max} = 0.988$

8036 measured reflections
 2160 independent reflections
 1722 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.118$
 $S = 1.05$
 2160 reflections
 110 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg_2 is the centroid of the C3–C8 phenyl ring.

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|---|--------------|--------------------|-------------|----------------------|
| $\text{N4}-\text{H2N4}\cdots\text{N1}^{\text{i}}$ | 0.871 (16) | 2.208 (16) | 3.0709 (18) | 171.1 (15) |
| $\text{C1}-\text{H1A}\cdots\text{N2}^{\text{ii}}$ | 0.93 | 2.50 | 3.4035 (16) | 166 |
| $\text{N4}-\text{H1N4}\cdots\text{Cg}_2^{\text{iii}}$ | 0.87 (2) | 2.58 (2) | 3.3929 (16) | 156.0 (17) |

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $x - 1, y, z$; (iii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and CKQ thank Universiti Sains Malaysia (USM) for the Research University Grant (No. 1001/PFIZIK/811160). CKQ also thanks USM for the award of a USM fellowship. AMI is thankful to the Director of the National Institute of Technology for providing research facilities and also thanks the Board for Research in Nuclear Sciences, Department of Atomic Energy, Government of India, for a Young Scientist Award.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5764).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fun, H.-K., Quah, C. K., Malladi, S. & Isloor, A. M. (2010). *Acta Cryst. E* **66**, o2799–o2800.
- Holla, B. S., Akberali, P. M. & Shivananda, M. K. (2000). *Farmaco*, **55**, 256–263.
- Isloor, A. M., Kalluraya, B. & Rao, M. (2000). *J. Saudi Chem. Soc.* **4**, 265–270.
- Isloor, A. M., Kalluraya, B. & Shetty, P. (2009). *Eur. J. Med. Chem.* **44**, 3784–3787.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Soliman, R., Habib, N. S., Ashour, F. A. & el-Taiebi, M. (2001). *Boll. Chim. Farm.* **140**, 140–148.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Sunil, D., Isloor, A. M. & Shetty, P. (2009). *Pharma Chem.* **1** 19–26.

supplementary materials

Acta Cryst. (2011). E67, o164 [doi:10.1107/S1600536810052025]

4-(1,2,4-Triazol-1-yl)aniline

H.-K. Fun, C. K. Quah, B. Chandrakantha, A. M. Isloor and P. Shetty

Comment

Compounds incorporating heterocyclic ring systems continue to attract considerable interests due to the wide range of biological activities they possess (Isloor *et al.*, 2000). Triazoles are a class of heterocyclic compounds having a five-membered ring of two carbon atoms and three nitrogen atoms (Soliman *et al.*, 2001). They have wide range of applications. In last few decades, triazoles have received much significant attention in the field of medicinal chemistry because of their diversified biological properties like antibacterial (Isloor *et al.*, 2009) and antifungal (Holla *et al.*, 2000) activities. In recent years, 1,2,4-triazole derivatives have been found to associate with anticancer properties (Sunil *et al.*, 2009). It is also observed that incorporation of aryl constituent into the triazoles ring systems augments the biological activities considerably.

In the title molecule (Fig. 1), the triazol-1-yl ring (N1-N3/C1/C2, maximum deviation = 0.003 (1) Å at atom N3) is inclined at angle of 34.57 (7)° with phenyl (C3-C8) ring. Bond lengths (Allen *et al.*, 1987) and angles are within normal ranges and are comparable to a related structure (Fun *et al.*, 2010).

In the crystal packing (Fig. 2), the molecules are linked into two-dimensional sheets parallel to the *ac*-plane via intermolecular N4–H2N4···N1 and C1–H1A···N2 hydrogen bonds. π - π stacking interactions between the centroids of N1-N3/C1/C2 triazol-1-yl rings (Cg1), with Cg1···Cg1ⁱ distance of 3.6750 (8) Å [symmetry code: (i) 1-X, -Y, 2-Z] are observed. The crystal structure is further consolidated by N4–H1N4···Cg2 (Table 1) interactions, where Cg2 is the centroid of C3-C8 phenyl ring.

Experimental

1,2,4-Triazole (2 g, 0.02 mol) was added lot-wise to a suspension of sodium hydride (60%, 1.47 g, 0.0308 mol) in dry DMF (20 ml) at 0°C. After the addition, the reaction mixture was stirred at the same temperature for 30 min. A solution of 4-fluoro nitrobenzene (2.82 g, 0.02 mol) in dry DMF (20 ml) was then added and the reaction mixture was stirred at room temperature for 18 h. The reaction mixture was then quenched with ice water and extracted with ethyl acetate. The organic layer was concentrated to afford a yellow solid as a nitro compound intermediate (3 g). This nitro compound was taken in methanol (30 ml) and hydrogenated using 10% palladium on carbon (0.2 g) at 3-kg pressure of hydrogen. After the reaction was over, the catalyst was filtered, the filtrate was concentrated to afford the title compound as a yellow solid. Yellow blocks were recrystallised from ethanol. Yield : 2.8g, 60 %. *M.p.* 433-435K.

Refinement

H1N4 and H2N4 were located in a difference Fourier map and allowed to refined freely. The remaining H atoms were positioned geometrically and refined using a riding model with C–H = 0.93 Å and $U_{iso}(H) = 1.2 U_{eq}(C)$. The highest residual electron density peak is located at 0.67 Å from C3 and the deepest hole is located at 1.27 Å from C8.

Figures




Fig. 1. The molecular structure of the title compound showing 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. The crystal structure of the title compound, viewed along the *b* axis. H atoms not involved in hydrogen bonds (dashed lines) have been omitted for clarity.

4-(1,2,4-Triazol-1-yl)aniline

Crystal data

$C_8H_8N_4$

$M_r = 160.18$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 5.5488$ (1) Å

$b = 7.3656$ (2) Å

$c = 19.5477$ (5) Å

$\beta = 99.416$ (2)°

$V = 788.15$ (3) Å³

$Z = 4$

$F(000) = 336$

$D_x = 1.350$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3245 reflections

$\theta = 3.0$ – 29.1 °

$\mu = 0.09$ mm⁻¹

$T = 296$ K

Block, yellow

$0.50 \times 0.42 \times 0.14$ mm

Data collection

Bruker SMART APEXII CCD diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2009)

$T_{\min} = 0.957$, $T_{\max} = 0.988$

8036 measured reflections

2160 independent reflections

1722 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 29.4$ °, $\theta_{\min} = 2.1$ °

$h = -7$ → 7

$k = -10$ → 10

$l = -26$ → 24

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.118$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0509P)^2 + 0.1531P]$

where $P = (F_o^2 + 2F_c^2)/3$

| | |
|--|---|
| $S = 1.05$ | $(\Delta/\sigma)_{\max} = 0.001$ |
| 2160 reflections | $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$ |
| 110 parameters | $\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$ |
| 0 restraints | Extinction correction: <i>SHELXTL</i> (Sheldrick, 2008), $F_c^* = kF_c[1+0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$ |
| Primary atom site location: structure-invariant direct methods | Extinction coefficient: 0.042 (5) |

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|--------------|--------------|-------------|----------------------------------|
| N1 | 0.3298 (2) | 0.17124 (18) | 1.08332 (6) | 0.0542 (3) |
| N2 | 0.58582 (19) | 0.24895 (17) | 1.00992 (6) | 0.0498 (3) |
| N3 | 0.34993 (17) | 0.26004 (14) | 0.97774 (5) | 0.0378 (3) |
| N4 | 0.0895 (3) | 0.4820 (2) | 0.70184 (6) | 0.0597 (4) |
| C1 | 0.2029 (2) | 0.2126 (2) | 1.02227 (7) | 0.0480 (3) |
| H1A | 0.0334 | 0.2092 | 1.0117 | 0.058* |
| C2 | 0.5611 (3) | 0.1960 (2) | 1.07255 (7) | 0.0533 (4) |
| H2A | 0.6948 | 0.1768 | 1.1072 | 0.064* |
| C3 | 0.2865 (2) | 0.31773 (16) | 0.90734 (6) | 0.0360 (3) |
| C4 | 0.4336 (2) | 0.27300 (17) | 0.85909 (6) | 0.0411 (3) |
| H4A | 0.5752 | 0.2056 | 0.8724 | 0.049* |
| C5 | 0.3691 (2) | 0.32893 (18) | 0.79107 (6) | 0.0432 (3) |
| H5A | 0.4686 | 0.2986 | 0.7589 | 0.052* |
| C6 | 0.1573 (2) | 0.43007 (17) | 0.76988 (6) | 0.0404 (3) |
| C7 | 0.0123 (2) | 0.47403 (17) | 0.81966 (6) | 0.0434 (3) |
| H7A | -0.1296 | 0.5415 | 0.8068 | 0.052* |
| C8 | 0.0763 (2) | 0.41891 (17) | 0.88743 (6) | 0.0415 (3) |
| H8A | -0.0218 | 0.4496 | 0.9200 | 0.050* |
| H2N4 | 0.171 (3) | 0.447 (2) | 0.6697 (8) | 0.065 (5)* |
| H1N4 | -0.029 (3) | 0.560 (3) | 0.6921 (9) | 0.078 (6)* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|------------|------------|------------|------------|------------|------------|
| N1 | 0.0549 (7) | 0.0658 (8) | 0.0435 (6) | 0.0049 (6) | 0.0124 (5) | 0.0063 (5) |

supplementary materials

| | | | | | | |
|----|------------|------------|------------|-------------|-------------|-------------|
| N2 | 0.0340 (5) | 0.0681 (8) | 0.0452 (6) | -0.0015 (5) | 0.0006 (4) | 0.0051 (5) |
| N3 | 0.0318 (5) | 0.0448 (5) | 0.0367 (5) | 0.0009 (4) | 0.0052 (4) | -0.0009 (4) |
| N4 | 0.0711 (8) | 0.0683 (9) | 0.0406 (6) | 0.0212 (7) | 0.0116 (6) | 0.0085 (6) |
| C1 | 0.0392 (6) | 0.0616 (8) | 0.0447 (7) | 0.0024 (6) | 0.0115 (5) | 0.0042 (6) |
| C2 | 0.0485 (7) | 0.0650 (9) | 0.0439 (7) | 0.0014 (6) | -0.0002 (6) | 0.0047 (6) |
| C3 | 0.0324 (5) | 0.0396 (6) | 0.0355 (6) | -0.0019 (4) | 0.0043 (4) | -0.0023 (4) |
| C4 | 0.0313 (5) | 0.0480 (7) | 0.0445 (7) | 0.0041 (5) | 0.0074 (5) | -0.0005 (5) |
| C5 | 0.0388 (6) | 0.0518 (7) | 0.0412 (6) | 0.0008 (5) | 0.0133 (5) | -0.0019 (5) |
| C6 | 0.0427 (6) | 0.0395 (6) | 0.0386 (6) | -0.0027 (5) | 0.0051 (5) | -0.0006 (5) |
| C7 | 0.0388 (6) | 0.0453 (7) | 0.0451 (7) | 0.0081 (5) | 0.0042 (5) | -0.0005 (5) |
| C8 | 0.0375 (6) | 0.0468 (7) | 0.0413 (6) | 0.0053 (5) | 0.0096 (5) | -0.0049 (5) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|--------------|--------------|-------------|--------------|
| N1—C1 | 1.3183 (17) | C3—C4 | 1.3842 (16) |
| N1—C2 | 1.3468 (19) | C3—C8 | 1.3851 (16) |
| N2—C2 | 1.3134 (18) | C4—C5 | 1.3821 (17) |
| N2—N3 | 1.3587 (14) | C4—H4A | 0.9300 |
| N3—C1 | 1.3332 (16) | C5—C6 | 1.3957 (17) |
| N3—C3 | 1.4284 (14) | C5—H5A | 0.9300 |
| N4—C6 | 1.3758 (16) | C6—C7 | 1.3987 (17) |
| N4—H2N4 | 0.871 (18) | C7—C8 | 1.3755 (16) |
| N4—H1N4 | 0.87 (2) | C7—H7A | 0.9300 |
| C1—H1A | 0.9300 | C8—H8A | 0.9300 |
| C2—H2A | 0.9300 | | |
| C1—N1—C2 | 102.04 (11) | C8—C3—N3 | 119.62 (10) |
| C2—N2—N3 | 102.11 (11) | C5—C4—C3 | 119.72 (11) |
| C1—N3—N2 | 109.16 (10) | C5—C4—H4A | 120.1 |
| C1—N3—C3 | 128.78 (10) | C3—C4—H4A | 120.1 |
| N2—N3—C3 | 122.06 (10) | C4—C5—C6 | 121.17 (11) |
| C6—N4—H2N4 | 121.5 (11) | C4—C5—H5A | 119.4 |
| C6—N4—H1N4 | 118.2 (12) | C6—C5—H5A | 119.4 |
| H2N4—N4—H1N4 | 120.1 (16) | N4—C6—C5 | 121.27 (12) |
| N1—C1—N3 | 111.01 (12) | N4—C6—C7 | 120.72 (12) |
| N1—C1—H1A | 124.5 | C5—C6—C7 | 118.00 (11) |
| N3—C1—H1A | 124.5 | C8—C7—C6 | 120.95 (11) |
| N2—C2—N1 | 115.69 (12) | C8—C7—H7A | 119.5 |
| N2—C2—H2A | 122.2 | C6—C7—H7A | 119.5 |
| N1—C2—H2A | 122.2 | C7—C8—C3 | 120.16 (11) |
| C4—C3—C8 | 120.00 (11) | C7—C8—H8A | 119.9 |
| C4—C3—N3 | 120.38 (10) | C3—C8—H8A | 119.9 |
| C2—N2—N3—C1 | -0.42 (15) | C8—C3—C4—C5 | -0.32 (19) |
| C2—N2—N3—C3 | 178.65 (11) | N3—C3—C4—C5 | 179.64 (11) |
| C2—N1—C1—N3 | -0.25 (16) | C3—C4—C5—C6 | 0.0 (2) |
| N2—N3—C1—N1 | 0.44 (16) | C4—C5—C6—N4 | -178.34 (13) |
| C3—N3—C1—N1 | -178.55 (12) | C4—C5—C6—C7 | 0.26 (19) |
| N3—N2—C2—N1 | 0.29 (17) | N4—C6—C7—C8 | 178.50 (13) |
| C1—N1—C2—N2 | -0.04 (18) | C5—C6—C7—C8 | -0.11 (19) |
| C1—N3—C3—C4 | -146.03 (13) | C6—C7—C8—C3 | -0.3 (2) |

| | | | |
|-------------|--------------|-------------|--------------|
| N2—N3—C3—C4 | 35.09 (17) | C4—C3—C8—C7 | 0.48 (19) |
| C1—N3—C3—C8 | 33.93 (19) | N3—C3—C8—C7 | -179.49 (11) |
| N2—N3—C3—C8 | -144.95 (12) | | |

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C3–C8 phenyl ring.

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|-------------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N4—H2N4 \cdots N1 ⁱ | 0.871 (16) | 2.208 (16) | 3.0709 (18) | 171.1 (15) |
| C1—H1A \cdots N2 ⁱⁱ | 0.93 | 2.50 | 3.4035 (16) | 166 |
| N4—H1N4 \cdots Cg2 ⁱⁱⁱ | 0.87 (2) | 2.58 (2) | 3.3929 (16) | 156.0 (17) |

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $x-1, y, z$; (iii) $-x, y+1/2, -z+3/2$.

Fig. 1




Fig. 2

